

# Metastatic Adenocarcinoma in the Brain



Ref.no.: MP2007

# Metastatic Adenocarcinoma in the Brain

## Clinical History

A 56-year old male underwent a total gastrectomy and splenectomy for gastric adenocarcinoma. Over a period of two months he developed a progressively unsteady gait, increasing weakness of his left hand and frontal headaches associated with nausea and vomiting. Imaging revealed a lesion in the right frontal lobe. He underwent a craniotomy with resection of the lesion, which was confirmed metastatic gastric adenocarcinoma. He experienced gradual increasing symptoms as well as jaundice, deteriorating consciousness and papilloedema from increased intracranial pressure. Repeat imaging revealed recurrence of the right frontal metastatic lesion as well as liver metastases. The patient died 9 months after his initial gastrectomy surgery.

## Pathology

This brain specimen is cut in the coronal plane. A circumscribed, variegated, pink-grey tumour is evident in the right frontal lobe. The tumour is involving the grey and white matter. Compression of the right lateral ventricle by the lesion is apparent with shift of the midline structures also seen.

## Further Information

Stomach cancer is one of the most common causes of cancer-related death worldwide. Risk factors include male gender, diet, smoking and chronic *Helicobacter pylori* infection. The most common sites for metastases of gastric adenocarcinoma are the liver, peritoneum, lungs and bones. Brain metastases are rare, occurring in <1% of cases. Isolated brain metastases are very uncommon with them being more commonly seen in disseminated disease and associated with a poor prognosis. Palliative treatment may include surgery, radiotherapy, steroid, chemotherapy or a combination thereof.

